

University of Groningen

Asymmetric Synthesis of 5-Substituted γ -Lactones and Butenolides via Nucleophilic Additions to Oxycarbenium Ions Derived from 5(R)-(Menthylloxy)-4(R)-(phenylsulfanyl)-2(3H)-dihydrofuranone

Oeveren, Arjan van; Feringa, Bernard

Published in:
The Journal of Organic Chemistry

DOI:
[10.1021/jo960078r](https://doi.org/10.1021/jo960078r)

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
1996

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Oeveren, A. V., & Feringa, B. (1996). Asymmetric Synthesis of 5-Substituted γ -Lactones and Butenolides via Nucleophilic Additions to Oxycarbenium Ions Derived from 5(R)-(Menthylloxy)-4(R)-(phenylsulfanyl)-2(3H)-dihydrofuranone. *The Journal of Organic Chemistry*, 61(9). <https://doi.org/10.1021/jo960078r>

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

The publication may also be distributed here under the terms of Article 25fa of the Dutch Copyright Act, indicated by the "Taverne" license. More information can be found on the University of Groningen website: <https://www.rug.nl/library/open-access/self-archiving-pure/taverne-amendment>.

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Asymmetric Synthesis of 5-Substituted γ -Lactones and Butenolides via Nucleophilic Additions to Oxycarbenium Ions Derived from 5(*R*)-(Menthylloxy)-4(*R*)-(phenylsulfanyl)-2(3*H*)-dihydrofuranone

Arjan van Oeveren and Ben L. Feringa*

Department of Organic and Molecular Inorganic Chemistry,
Groningen Center for Catalysis and Synthesis,
University of Groningen, Nijenborgh 4,
9747 AG Groningen, The Netherlands

Received January 12, 1996

Optically active 5-alkyl-substituted butenolides and γ -lactones are attractive building blocks in natural product synthesis¹ and comprise structural moieties frequently present in, e.g., insect pheromones,² cardenolides,³ lignans, and flavor components.⁴ Efficient and stereoselective synthetic routes to these products in enantiomerically pure form are highly desirable.⁵ As part of our explorative studies toward the use of 5(*R*)-(menthylloxy)-2(5*H*)-furanone (**1**) as a chiral synthon,^{6,7} the enantioselective synthesis of a number of naturally occurring lignans has been reported.⁸

5(*R*)-(Menthylloxy)-2(5*H*)-furanone (**1**) reacts with thiophenol and a catalytic amount of triethylamine to give stereospecifically and in excellent yield the trans addition product **2** (Scheme 1), which features an attractive functional group arrangement to generate α -sulfanyl oxycarbenium ion **3**. Here we report the fast and highly stereoselective transformations of **2** to 5-alkyl-substituted 4-(phenylsulfanyl)-2(3*H*)-dihydrofuranones **4** or 4-(menthylloxy)-3-(phenylsulfanyl) carboxylic acids **5**, which are precursors for butenolides and γ -lactones.

Various methods for C–C bond formation via Lewis acid-mediated reactions of acetals with nucleophiles have been developed.⁹ As demonstrated by a number of groups,¹⁰ there is a mechanistic divergence between S_N2 -





Figure 1.



and S_N1 -type processes. It has been shown that Lewis acid-mediated additions of silylated nucleophiles to α - and β -sulfanyl-substituted aldehydes proceed with excellent diastereoselectivities.¹¹ Furthermore, reactions of α -sulfanyl acetals with carbon nucleophiles have been studied by Saigo and co-workers.¹² In furanone **2**, an α -sulfanyl-substituted, mixed acyloxy–alkoxy acetal moiety is present, and upon treatment with a Lewis acid it is observed that the acyloxy acetal bond is always broken¹³ and this reaction path leaves only two likely intermediates (Figure 1). The stereoselectivity of these reactions can be rationalized by the Felkin–Ahn model, and conformers **A** and **B** lead to anti and syn adducts, respectively. Conformer **A** is preferred, and in most cases the anti adduct is the only detectable diastereomer.

When 4(*R*)-(phenylsulfanyl)-substituted furanone **2** is treated at -70°C with 1–2 equiv of TiCl_4 in the presence of a variety of nucleophiles such as allylsilanes, silyl enol ethers, or diorganozinc reagents a very fast reaction occurs. After a reaction time of 5 min and subsequent aqueous workup, β , γ -substituted acids **5** are isolated in 56–72% yield and in most cases with a diastereomeric ratio $>98:2$ according to ^1H and ^{13}C NMR (Scheme 2, path A, Table 1, entries 2, 4, 6, 8, and 12). In addition to the acids **5** small amounts of lactones **4** (5–20%) are also found in the crude product, but these could be easily separated.

When the addition of nucleophiles **9**–**16** is performed in the presence of 2 equiv of TiCl_4 for 1 min at ambient temperature, followed by aqueous workup, the lactones **4** are obtained. In a few cases small amounts of the acids **5** ($\leq 10\%$) are also formed. Apparently, the intermediate **6** is activated by the excess of Lewis acid present, and upon addition of water hydrolysis to the hydroxy acid **8**

(1) (a) Hanessian S. In *The total synthesis of natural products: The chiron approach*; Pergamon Press: New York, 1983; Chapter 9. (b) Fernandez, A.-M.; Jacob, M.; Gralak, J.; Al-Bayati, Y.; Plé, G.; Duhamel, L. *Synlett* **1995**, 431. (c) Takahata, H.; Uchida, Y.; Momose, T. *J. Org. Chem.* **1994**, *59*, 7201. (d) Canan Koch, S. S.; Chamberlin, A. R. *J. Org. Chem.* **1993**, *58*, 2725.

(2) (a) Fukusaki, E.; Senda, S.; Nakazono, Y.; Omata, T. *Tetrahedron* **1991**, *47*, 6223. (b) Wheeler, J. W.; Happ, G. M.; Araujo, J.; Pasteels, J. M. *Tetrahedron Lett.* **1972**, *46*, 4635. (c) Mori, K. *Tetrahedron* **1989**, *45*, 3233.

(3) Rao, Y. S. *Chem. Rev.* **1976**, *78*, 625.

(4) (a) Bloch, R.; Gilbert, L. *J. Org. Chem.* **1987**, *52*, 4603. (b) Thijs, L.; Waanders, W. P.; Stokkingreef, E. H. M.; Zwanenburg, B. *Recl. Trav. Chim. Pays-Bas* **1986**, *105*, 332.

(5) For recent examples, see: (a) Brown, H. C.; Kulkarni, S. V.; Racherla, U. S. *J. Org. Chem.* **1994**, *59*, 365. (b) Miller, M.; Hegedus, L. S. *J. Org. Chem.* **1993**, *58*, 6779. (c) Doyle, M. P.; Protopenova, M. N.; Zhou, Q.-L.; Bode, J. W. *J. Org. Chem.* **1995**, *60*, 6654.

(6) For reviews see: (a) Feringa, B. L.; De Jong, J. C. *Bull. Soc. Chim. Belg.* **1992**, *101*, 627. (b) Feringa, B. L.; De Lange, B.; De Jong, J. C.; Lubben, M.; Faber, W.; Schudde, E. P. *Pure Appl. Chem.* **1992**, *64*, 1865. (c) De Jong, J. C.; Van Bolhuis, F.; Feringa, B. L. *Tetrahedron: Asymmetry* **1991**, *2*, 1247.

(7) (a) Feringa, B. L.; De Lange, B.; De Jong, J. C. *J. Org. Chem.* **1989**, *54*, 2471. (b) Rispens, M. T.; Keller, E.; De Lange, B.; Zijlstra, W. J.; Feringa, B. L. *Tetrahedron: Asymmetry* **1994**, *5*, 607. (c) Hulst, R.; De Vries, N. K.; Feringa, B. L. *J. Org. Chem.* **1994**, *59*, 7453.

(8) (a) Van Oeveren, A.; Jansen, J. F. G. A.; Feringa, B. L. *J. Org. Chem.* **1994**, *59*, 5999 and references cited. (b) Middel, O.; Woerdenbag, H. J.; Van Uden, W.; Van Oeveren, A.; Jansen, J. F. G. A.; Feringa, B. L.; Konings, A. T.; Pras, N.; Kellogg, R. M. *J. Med. Chem.* **1995**, *38*, 2112.

(9) For representative reviews, see: (a) Alexakis, A.; Mangeney, P. *Tetrahedron: Asymmetry* **1990**, *1*, 477. (b) Mukaiyama, T.; Murakami, M. *Synthesis* **1987**, 1043.


(10) (a) Mori, I.; Ishihara, K.; Flippin, L. A.; Nozaki, K.; Yamamoto, H.; Bartlett, P. A.; Heathcock, C. H. *J. Org. Chem.* **1990**, *55*, 6107. (b) Denmark, S. E.; Almstead, N. G. *J. Am. Chem. Soc.* **1991**, *113*, 8089. (c) Sammiaka, T.; Smith, R. S. *J. Am. Chem. Soc.* **1994**, *116*, 7915.

(11) Annunziata, R.; Cinquini, M.; Cozzi, F.; Cozzi, P. G.; Consolandi, E. *J. Org. Chem.* **1992**, *57*, 456.

(12) Kudo, K.; Hashimoto, Y.; Sukegawa, M.; Hasegawa, M.; Saigo, K. *J. Org. Chem.* **1993**, *58*, 579.

(13) Van Oeveren, A.; Feringa, B. L. *Tetrahedron Lett.* **1994**, *35*, 8437.


Scheme 2

Table 1. Additions of Nucleophiles to Oxycarbenium Ions Derived from **2**

Entry	Nucleophile	T ($^\circ\text{C}$)	time (min)	Product (R)	(ratio a)	Yield b %	trans : cis c (anti : syn c)
1		23	1	4a/5a	(>9 : 1)	50	92 : 8
2		-70	5	5a/4a	(8 : 2)	56	9 : 1
3		23	1	4b/5b	(>9 : 1)	65	>98 : 2
4		-70	5	5b/4b	(8 : 2)	59	>98 : 2
5		23	1	4c/5c	(>9 : 1)	66	>98 : 2
6		-70	5	5c/4c	(9 : 1)	72	>98 : 2
7		23	1	4d/5d	(>9 : 1)	74	>98 : 2 ^d
8		-70	5	5d/4d	(8 : 2)	52	>98 : 2 ^d
9		23	1	4e/5e	(>9 : 1)	57	>98 : 2 ^d
10		23	1	4f/5f	(>9 : 1)	74	53 : 47 ^d
11		23	2	4g/5g	(>9 : 1)	56	9 : 1
12		-70	5	5g/4g	(9 : 1)	57	>98 : 2
13		23	5	4h/5h	(>9 : 1)	53	>98 : 2

^aRatio determined by ^1H NMR. ^bIsolated yield of the major isomer.

^cRatio determined by ^1H and ^{13}C NMR of the crude product. ^dDiastereomeric ratio of the exocyclic stereogenic center was in the order of 6 : 4 in all four cases (entry 7-10).




followed by lactonization takes place (Scheme 2, path B). A key feature of the second Lewis acid-mediated step is the stereoselective cleavage of the auxiliary group assisted by the α -sulfanyl functionality. Except for **4f**, trans lactones **4** are formed as the major, or in most cases only detectable, diastereomer. Even in the few cases (entries 1 and 11, Table 1) where small amounts of cis lactones **4** were obtained, these could be separated by simple column chromatography using silica gel.


Single crystal X-ray analysis of **4b** confirmed the trans relationship of the substituents at C_4 and C_5 .²¹ The trans configuration for the products **4a-e,g,h** has been assigned by comparison on the basis of ^1H NMR coupling constants and on the expected mechanistic similarity in this reaction for all nucleophiles used.¹⁴

The enantiomerically pure 5(*S*)-alkyl-4(*R*)-(phenylsulfanyl)-2(3*H*)-dihydrofuranones **4** are excellent precursors for 5-alkyl-substituted butenolides or butanolides,^{15,16} whereas the phenylsulfanyl substituent allows a number

Scheme 3



Scheme 4



of synthetically useful transformations. Reductive desulfurization of **4** could best be performed with Bu_3SnH /AIBN and provided enantiomerically pure 5(*S*)-alkyl-2(3*H*)-dihydrofuranones **17** (Scheme 3). The flavor components and insect pheromones⁴ 5(*S*)-ethyl-2(3*H*)-dihydrofuranone (**17g**) [85% yield, $[\alpha]_D = -50$ (c 1, MeOH) (lit.¹⁷ $[\alpha]_D = -53.2$ (c 1, MeOH))] and 5(*S*)-octyl-2(3*H*)-dihydrofuranone (**17h**) [66% yield, $[\alpha]_D = -35$ (c 0.48, MeOH) (lit.¹⁷ $[\alpha]_D = -36.8$ (c 0.3, MeOH))] were obtained using this procedure.

Alternatively, 5(*S*)-alkyl-2(5*H*)-furanones **19** are accessible via the sulfones **18** (Scheme 4). Oxidation of **4** with Oxone¹⁸ and subsequent elimination with basic alumina in CH_2Cl_2 gave the corresponding 5(*S*)-alkyl-2(5*H*)-furanones **19** in good yields.¹⁹ For example, 5(*S*)-ethyl-2(5*H*)-furanone (**19g**) [95% overall yield, $[\alpha]_D = +105$ (c 4.1, CH_2Cl_2) (lit.²⁰ for 5(*R*)-**19g** $[\alpha]_D = -97.6$ (c 2.08, CH_2Cl_2))] and 5(*S*)-(1-prop-2-enyl)-2(5*H*)-furanone (**19a**) (72% overall yield, $[\alpha]_D = +105$ (c 1.08, MeOH)) were obtained using this procedure.

In conclusion, we have demonstrated that **2** is a valuable synthon for the synthesis of 3,4-disubstituted carboxylic acids, 5-substituted butenolides, and γ -lactones in enantiomerically pure form.

Acknowledgment. Financial support was provided by the Netherlands Organization for Scientific Research (NWO/SOON).

Supporting Information Available: Experimental procedures for the Lewis acid mediated additions and spectroscopic data for compounds **4a-h** and **5a,b,c,d,g**. (5 pages).

JO960078R

(16) Carretero, J. C.; De Lombaert, S.; Ghosez, L. *Tetrahedron Lett.* **1987**, 28, 2135.

(17) Roder, H.; Helmchen, G.; Peters, E.-M.; Peters, K.; Von Schnering, H.-G. *Angew. Chem., Int. Ed. Engl.* **1984**, 23, 898.

(18) Trost, B. M.; Curran, D. P. *Tetrahedron Lett.* **1981**, 22, 1287.

(19) Vidal, J.; Huet, F. *Tetrahedron Lett.* **1986**, 27, 3733.

(20) Cardellach, J.; Font, J.; Ortuño, R. M. *J. Heterocyclic Chem.* **1984**, 21, 327.

(21) The author has deposited atomic coordinates for **4b** with the Cambridge Crystallographic Data Centre. The coordinates can be obtained, on request, from the Director, Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge, CB2 1EZ, UK.

(14) See the supporting information.

(15) López, C.; Gómez, A. M.; Valverde, S. *Synlett* **1991**, 825.